

Written on 12 May 2025 3 minutes of reading  
News

- Fundamental Research
- [Geosciences](#)
- [Numerical methods and optimization](#)

**The climate is changing and it is doing so too fast. The impacts on our environment are already visible in many regions of the world. In such a context, it is important to understand the dynamics of our environment in order to construct tools capable of anticipating the risks stemming from this climate disruption. Among these tools, numerical models offer a very effective solution for testing a wide range of climate scenarios, taking into account numerous physical, chemical and biological processes.**

**For example, numerical landscape evolution models are widely used in order to understand and quantify the relationship between the climate and the hydro-sedimentary dynamics of a water catchment area.** Two ingredients are required to construct models of this type: a “good ladleful” of a water flow model and a “large cup” of sedimentary transport, all bound together by the pooled expertise of geoscientists and applied mathematicians! This recipe formed the foundation for the Dionisos software solution [1], which became [DionisosFlow](#) [2], designed for the purposes of modeling hydro-sedimentary processes.

### **A base ingredient: “numerical consistency”**

One of the “sensitive” aspects of these models, which have been known for over 20 years, is a much greater dependence on mesh size and shape<sup>1</sup> than can usually be attributed to the simple numerical discretization of our continuous equations. For a long time, the origin of this dependence remained unclear. In 2020, a first article [2] was published identifying one of the reasons for which some water flow models conventionally used by the community produced an artificial mesh dependence. The reason turned out to be simple: the flow algorithm, which was supposed to mimic an “empirical” model, could actually be traced back established physics, namely a Manning-Strickler model<sup>2</sup>. However, these algorithms had not been explicitly designed to reproduce established physics, expressible through a set of partial differential equations. And since chance does not always get things quite right, it turns out that the algorithms required a few tweaks to ensure this was the case. These tweaks are referred to as “numerical consistency”. Without it, an algorithm is bereft of its physics: it still models something, but something completely artificial! This defect, present in historical models, can now be corrected for most embedded algorithms [3].

<sup>1</sup> A numerical domain being made up of a set of elementary meshes

<sup>2</sup> An empirical formula used to estimate the average velocity of a liquid flowing in a conduit with [a free surface](#), i.e., flowing in a conduit that does not completely enclose the fluid or in an open channel.

### **An essential utensil: a filter to remove lumps**

However, solving this problem is not enough to avoid “lumps” (those tiny flaws in the numerical solution which, when amplified, spoil the dish!) in landscape evolution models. The other aspect that needs to be considered is the coupling between water flow and sediment transport. To do this, we adapted an approach commonly deployed in LES<sup>3</sup>-type models to eliminate the “lumps” arising from turbulence below a certain size [4]. Concretely, this means that, on the one hand, water flow is governed by a filtered topography, and on the other hand, only sediments transported through a similarly filtered hydrological network contribute to shaping the topography. The key ingredient is known as the Leray-alpha<sup>4</sup> filter: in addition to fulfilling its intended function, this approach preserves the non-linear effects arising from flow–transport coupling, while effectively filtering out numerical noise<sup>5</sup>.

<sup>3</sup> Large Eddy Simulation : méthode de modélisation de la turbulence utilisée dans le domaine de la dynamique des fluides

<sup>4</sup> Regularization method by local averaging of alpha size

<sup>5</sup> Solution errors associated with the numerical resolution process

## Reproducible results from one simulation to the next

The benefits of such a method became clear when it was realized that the reproducibility of results was not guaranteed in its absence. This occurred, for example, when the problem was solved with different solvers for the same simulation, or indeed simply when the number of processors involved in solving the numerical system was varied. The results obtained were then considerably altered (a river, instead of flowing straight ahead, could randomly fork), but this has now been resolved thanks to the use of the filter (Figure 1).




Figure 1: numerical results obtained without and with the use of the filter. The three simulations are strictly similar in their physical parameters but the systems of equations are solved using different solvers.

**With the right proportions, the dish is ready**

As with any good recipe, it is important to get the proportions right. And the same is true here, since the size of the filter to be chosen is governed by the size of the objects we want to preserve in our landscape. An object such as a mountain, a drainage basin or even a simple river should be approximately one order of magnitude larger than the size of the filter. As long as this rule is followed, all that's left to do is adapt the mesh size so that it can “capture” the filter correctly, and it’s ready!(Figure 2)




Figure 2 : Example of a complex simulation with filter and realistic physics. In this simulation, the topography is carved out by the flow of water and valleys develop

## References :

- [1] D. Granjeon (1996). *Modélisation stratigraphique déterministe : Conception et applications d'un modèle diffusif 3-d multi-lithologique.*(*Deterministic stratigraphic modeling: Design and applications of a 3d multi-lithology diffusion model*) PhD thesis, Université de Rennes I
- [2] J. Coatléven (2020). *Some multiple flow direction algorithms for overland flow on general meshes*, ESAIM: Mathematical Modelling and Numerical Analysis.  
>> DOI: <https://doi.org/10.1051/m2an/2020025>
- [3] J. Coatléven; B. Chauveau (2025). *A post-processing solution to restore numerical consistency for classical flow routing algorithms. Computational Geosciences*  
>> DOI : <https://doi.org/10.1007/s10596-025-10359-5>
- [4] J. Coatléven; B. Chauveau (2024). *Large structure simulation for landscape evolution models, Earth Surface Dynamics.*  
>> DOI: <https://doi.org/10.5194/esurf-12-995-2024>

Scientific contacts : [Benoît Chauveau](#), [Julien Coatléven](#)

## You may also be interested in

[Global change, impact on landscapes and water resources](#)

[Lacustrine sedimentary series: an archive of past environmental changes to better understand the present](#)

[Knowledge and description of sedimentary systems](#)

A numerical recipe for adapting the landscape map

12 May 2025

Link to the web page :